

UNITED STATES ATTORNEY'S OFFICE FOR THE DISTRICT OF HAWAII

PROTECT YOURSELF AGAINST CORONAVIRUS SCAMS

Crooks are taking advantage of the current pandemic to defraud seniors. Here are tips to help you avoid being victimized by scammers.

DON'T SHARE YOUR BANKING OR PERSONAL INFORMATION OVER THE PHONE. The IRS will never ask for this information via telephone.

DON'T CLICK ON EMAIL LINKS FROM UNKNOWN SOURCES. Links from unknown senders could contain harmful viruses.

DON'T BUY FROM AN UNFAMILIAR ONLINE BUSINESS. Avoid buying supplies like gloves, masks, etc., from online sellers you are not familiar with.

DO HANG UP ON ROBOCALLS. If you answer the phone and hear a pre-recorded voice, be careful of scam offers and requests.

DO VERIFY YOUR BILL COLLECTOR'S IDENTITY. If you must make payments via telephone, only do so by calling the phone number on your bill.

DO CHECK BEFORE YOU GIVE. Make sure the charity of your choice is legitimate before you give them payment information.



REPORT SCAMS TO:

UNITED STATES ATTORNEY'S OFFICE FOR THE DISTRICT OF HAWAII:

usahi.covid19@usdoj.gov T: 1.808.541.2850

HAWAII STATE OFFICE OF CONSUMER PROTECTION:

cca.hawaii.gov/ocp/ consumer-complaint/ T: 1.808.587.4272

NATIONAL CENTER FOR DISASTER FRAUD:

disaster@leo.gov T: 1.866.720.5721

ADDITIONAL INFO:

www.cdc.gov

www.coronavirus.gov

ftc.gov/coronavirus

www.justice.gov/usaohi/covid-19-fraud

SEQUENCE FOR PUTTING ON PERSONAL PROTECTIVE EQUIPMENT (PPE)

The type of PPE used will vary based on the level of precautions required, such as standard and contact, droplet or airborne infaction isolation precautions. The procedure for putting on and removing PPE should be tailored to the specific type of PPE.

1. GOWN

- Fully cover torso from neck to knees, arms to end of wrists, and wrap around the back
- Fasten in back of neck and waist

2. MASK OR RESPIRATOR

- Secure ties or elastic bands at middle of head and neck
- · Fit flexible band to nose bridge
- · Fit snug to face and below chin
- Fit-check respirator

3. GOGGLES OR FACE SHIELD

· Place over face and eyes and adjust to fit



4. GLOVES


· Extend to cover wrist of isolation gown

USE SAFE WORK PRACTICES TO PROTECT YOURSELF AND LIMIT THE SPREAD OF CONTAMINATION

- · Keep hands away from face
- · Limit surfaces touched
- · Change gloves when torn or heavily contaminated
- · Perform hand hygiene



042306123



HOW TO SAFELY REMOVE PERSONAL PROTECTIVE EQUIPMENT (PPE) EXAMPLE 1

There are a variety of ways to safely remove PPE without contaminating your clothing, skin, or mucous membranes with potentially infectious materials. Here is one example. Remove all PPE before exiting the patient room except a respirator, if worn. Remove the respirator after leaving the patient room and closing the door. Remove PPE in the following sequence:

1. GLOVES

- · Outside of gloves are contaminated!
- If your hands get contaminated during glove removal, immediately wash your hands or use an alcohol-based hand sanitizer
- Using a glowed hand, grasp the palm area of the other glowed hand and peel off first glowe
- · Hold removed glove in gloved hand
- Slide fingers of ungloved hand under remaining glove at wrist and peel off second glove over first glove
- Discard gloves in a waste container

2. GOGGLES OR FACE SHIELD

- Outside of goggles or face shield are contaminated!
 If your hands get contaminated during goggle or face shield removal,
- If your hands get contaminated during goggle or face shield removal, immediately wash your hands or use an alcohol-based hand sanitizer
 Remove goggles or face shield from the back by lifting head band or
- ear pieces

 If the item is reusable, place in designated receptacle for
- reprocessing. Otherwise, discard in a waste container

3. GOWN

- Gown front and sleeves are contaminated!
 If your hands get contaminated during gown removal, immediately wash your hands or use an alcohol-based hand sanitizer
- Unfasten gown ties, taking care that sleeves don't contact your body when reaching for ties
- when reaching for the
- Pull gown away from neck and shoulders, touching inside of gown only
- Turn gown inside out

Fold or roll into a bundle and discard in a waste container

4. MASK OR RESPIRATOR

- Front of mask/respirator is contaminated DO NOT TOUCH!
- If your hands get contaminated during mask/respirator removal, immediately wash your hands or use an alcohol-based hand sanitizer
- Grasp bottom ties or elastics of the mask/respirator, then the ones at
- the top, and remove without touching the front
 Discard in a waste container
- 5. WASH HANDS OR USE AN ALCOHOL-BASED HAND SANITIZER IMMEDIATELY AFTER REMOVING ALL PPE

PERFORM HAND HYGIENE BETWEEN STEPS IF HANDS BECOME CONTAMINATED AND IMMEDIATELY AFTER REMOVING ALL PPE







HOW TO SAFELY REMOVE PERSONAL PROTECTIVE EQUIPMENT (PPE) EXAMPLE 2

Here is another way to safely remove PPE without contaminating your clothing, skin, or mucous membranes with potentially infectious materials. Remove all PPE before exiting the patient room except a respirator, if worn. Remove the respirator after leaving the patient room and closing the door. Remove PPE in the following sequence:

1. GOWN AND GLOVES

- Gown front and sleeves and the outside of gloves are contaminated!
- If your hands get contaminated during gown or glove removal, immediately wash your hands or use an alcohol-based hand sanitizer
- Grasp the gown in the front and pull away from your body so that the ties break, touching outside of gown only with gloved hands
- While removing the gown, fold or roll the gown inside-out into a bundle
- As you are removing the gown, peel off your gloves at the same time, only touching the inside of the gloves and gown with your bare hands. Place the gown and gloves into a waste container.



OR

2. GOGGLES OR FACE SHIELD

- Outside of goggles or face shield are contaminated!
- If your hands get contaminated during goggle or face shield removal, immediately wash your hands or use an alcohol-based hand senitizer
 Remove goggles or face shield from the back by lifting head band and
- Remove goggles or face shield from the back by litting head band a without touching the front of the goggles or face shield
 If the item is reusable, place in designated receptacle for
- If the item is reusable, place in designated receptacle to reprocessing. Otherwise, discard in a waste container

3. MASK OR RESPIRATOR

- Front of mask/respirator is contaminated D0 NDT T0UCH!
 If your hands get contaminated during mask/respirator removal,
- If your hands get contaminated during mask/respirator removal, immediately wash your hands or use an alcohol-based hand sanitizer
 Grasp bottom ties or elastics of the mask/respirator, then the ones at
- Grasp bottom deside ensues of ensues of the mask respirator, men de creatities the top, and remove without touching the front
 Origonal in a work operations
- Discard in a waste container

4. WASH HANDS OR USE AN ALCOHOL-BASED HAND SANITIZER IMMEDIATELY AFTER REMOVING ALL PPE

PERFORM HAND HYGIENE BETWEEN STEPS IF HANDS BECOME CONTAMINATED AND IMMEDIATELY AFTER REMOVING ALL PPE



Use Personal Protective Equipment (PPE) When Caring for Patients with Confirmed or Suspected COVID-19

Before caring for patients with confirmed or suspected COVID-19, healthcare personnel (HCP) must:

- Receive comprehensive training on when and what PPE is necessary, how to don (put on) and doff (take off) PPE, limitations
 of PPE, and proper care, maintenance, and disposal of PPE.
- Demonstrate competency in performing appropriate infection control practices and procedures.

Remember:

- + PPE must be donned correctly before entering the patient area (e.g., isolation room, unit if cohorting).
- PPE must remain in place and be worn correctly for the duration of work in potentially contaminated areas. PPE should not be adjusted (e.g., retying gown, adjusting respirator/facemask) during patient care.
- FPE must be removed slowly and deliberately in a sequence that prevents self-contamination. A step-by-step process should be developed and used during training and patient care.



CONTRA MINISTER

Donning (putting on the gear):

More than one domning method may be acceptable. Training and practice using your healthcare facility's procedure is critical. Below is one example of domning.

- 1. Identify and gather the proper PPE to don. Ensure choice of gown size is correct (based on training).
- 2. Perform hand hygiene using hand sanitizer.
- 3. Put on isolation gown. The all of the ties on the gown. Assistance may be needed by another HCP.
- 4. Put on NIOSH-approved N95 filtering facepiece respirator or higher (use a facemask if a respirator is not available). If the respirator has a nosepiece, it should be fitted to the nose with both hands, not bent or tented. Do not pinch the nosepiece with one hand. Respirator/facemask should be extended under chin. Both your mouth and nose should be protected. Do not wear respirator/facemask under your chin or store in scrubs pocket between patients."
 - > Respirator: Respirator straps should be placed on crown of head (top strap) and base of neck (bottom strap). Perform a user seal check each time you put on the respirator.
 - Facemask: Mask ties should be secured on crown of head (top tie) and base of neck (bottom tie). If mask has loops, hook them appropriately around your ears.
- 5. Put on face shield or goggles. When wearing an N95 respirator or half facepiece elastomeric respirator, select the proper eye protection to ensure that the respirator does not interfere with the correct positioning of the eye protection, and the eye protection does not affect the fit or seal of the respirator. Face shields provide full face coverage. Goggles also provide excellent protection for eyes, but fogging is common.
- 6. Put on gloves. Gloves should cover the cuff (wrist) of gown.
- 7. HCP may now enter patient room.

Doffing (taking off the gear):

More than one doffing method may be acceptable. Training and practice using your healthcare facility's procedure is critical. Below is one example of doffing.

- Remove gloves. Ensure glove removal does not cause additional contamination of hands. Gloves can be removed using more than one technique (e.g., glove-in-glove or bird bealc).
- Remove gown. Untie all ties (or unsnap all buttons). Some gown ties can be broken rather than untied. Do so in gentle
 manner, avoiding a forceful movement. Reach up to the shoulders and carefully pull gown down and away from the body.
 Rolling the gown down is an acceptable approach. Dispose in trach receptacle.⁴
- 3. HCP may now exit patient room.
- 4. Perform hand hygiene.
- Remove face shield or goggles. Carefully remove face shield or goggles by grabbing the strap and pulling upwards and away from head. Do not touch the front of face shield or goggles.
- Remove and discard respirator (or facemask if used instead of respirator).* Do not touch the front of the respirator or facemask.
 - > Respirator: Remove the bottom strap by touching only the strap and bring it carefully over the head. Grasp the top strap and bring it carefully over the head, and then pull the respirator away from the face without touching the front of the respirator.
 - » Facemask: Carefully untie (or unbook from the ears) and pull away from face without touching the front.
- Perform hand hygiene after removing the respirator/facemask and before putting it on again if your workplace is practicing reuse.

"Facilities implementing reuse or extended use of PPE will need to adjust their donning and doffing procedures to accommodate those practices.

www.cdc.gov/coronavirus

Mahalo,

Angel England, RN Operations Manager Update Subscription Preferences Subscribe Forward to a Friend Mailing address: 500 Ala Moana Blvd, Suite 7400, Honolulu, HI 96813 Phone number: 808-234-5380

 This email was sent to << Email Address>>

 why did I get this?
 unsubscribe from this list
 update subscription preferences

 Community Ties of America, Inc. · 500 Ala Moana Blvd Suite 7400 · Honolulu, Hawaii 96813 · USA

